

SPECIMEN COLLECTION INSTRUCTIONS

Incyte Swab Specimens

Infectious Disease Testing

Incyte Diagnostics offers infectious disease testing using the Incyte swab. These gynecologic tests aid in the detection, diagnosis, and treatment of viral, fungal, and bacterial infections. Nucleic acid testing using Polymerase Chain Reaction (PCR) methodology is employed to provide high levels of sensitivity and specificity in testing for the presence of these organisms. The following tests are orderable using an Incyte swab collection.



- Chlamydia
- Gonorrhea
- Trichomonas vaginalis
- Gardnerella vaginalis
- Candida albicans
- Candida glabrata
- Candida tropicalis
- Candida parapsilosis
- Mycoplasma genitalium
- Mycoplasma hominis
- Atopobium vaginae
- Ureaplasma spp.
- Mobiluncus mulieris
- Mobiluncus curtisii

For patients with suspected Vaginosis/Vaginitis:

Please use Incyte swab for specimen collection

- A thorough collection of the vaginal walls is required.

For patients with Cervicitis:

Please use Incyte swab for specimen collection

- Begin with an initial thorough collection from the cervix. A subsequent vaginal collection may be made with the same swab if vaginal organism testing is also desired.

Specimen Collection:

- Aseptically remove sterile swab from package.
- Collect specimen by vigorously swabbing site for 30 seconds.
- Aseptically remove cap from vial.
- Place swab in transport medium and break off swab against rim of the tube.
- Replace cap to vial, close tightly.
- Fill out vial label with full patient name and one other identifier.

SPECIMEN COLLECTION INSTRUCTIONS


ThinPrep® Pap Specimen

Option 1

ThinPrep® sample collection with mop/broom device



1. Record the patient's name and ID number on the vial, and the patient information and medical history on the cytology requisition form.



2. Obtain an adequate sample from the transformation zone of the cervix using a cervical sampler (broom-like device). Insert the central bristles of the broom into the endocervical canal deep enough to allow the shorter bristles to fully contact the ectocervix. Push gently, and rotate the brush 360° in a clockwise direction 3-5 times.



3. Rinse the cervical sampler immediately into the vial by pushing it into the bottom of the vial 10 times, forcing the bristles apart. As a final step, swirl the brush **vigorously** to further release material. Visually inspect the cervical sampler to ensure that no material remains attached. Discard the collection device. **Do not leave the head of the cervical sampler in the vial.**




4. Tighten the cap so that the black torque line on the cap passes the black torque line on the vial. **Do not over-tighten.** Place the vial and pathology request form in a specimen bag for transportation to the laboratory.

Option 2

ThinPrep® sample collection with brush/spatula




1. Record the patient's name and ID number on the vial, and the patient information and medical history on the cytology requisition form.



2. Obtain an adequate sample from the ectocervix using a plastic spatula. Select the contoured end of the plastic spatula and rotate it 360° in a clockwise direction around the entire ectocervix, while maintaining tight contact with the ectocervical surface.



3. Rinse the spatula as quickly as possible into the vial by swirling the spatula **vigorously** in the vial 10 times. Discard the plastic spatula.



4. Obtain an adequate sampling from the endocervix using an endocervical brush device. Insert the Cytobrush into the endocervical canal until only the bottom-most bristles are exposed. Slowly rotate 1/4 or 1/2 turn in one direction. **Do not over rotate the brush.**



5. Rinse the endocervical brush immediately in the same vial by rotating the device in the solution 10 times while pushing it against the vial wall. As a final step, swirl the brush **vigorously** to further release material. Discard the brush.



6. Tighten the cap so that the torque line on the cap passes the torque line on the vial. **Do not over-tighten.** Place the vial and pathology request form in a specimen bag for transport to the laboratory.

Aptima® Multitest Swab Specimen Collection Kit

Clinician collection procedure guide


Collection for vaginal swab specimens




Partially open the swab package and remove the swab. Do not touch the soft tip or lay the swab down. **If the soft tip is touched, laid down, or dropped, discard and get a new Aptima Multitest Swab Specimen Collection Kit.** Hold the swab, placing thumb and forefinger in the middle of the shaft covering the black score line. Do not hold the shaft below the score line.

Swab specimen collection guide for:


- *Chlamydia trachomatis* (CT)
- *Neisseria gonorrhoeae* (NG)
- *Trichomonas vaginalis* (TV)



Carefully insert the swab into the vagina about 2 inches (5 cm) past the introitus and gently rotate the swab for 10 to 30 seconds. Make sure the swab touches the vaginal walls so that moisture is absorbed by the swab. Withdraw the swab without touching the skin.



While holding the swab in hand, unscrew the tube cap. Do not spill the tube contents. **If the tube contents are spilled, discard and replace with a new Aptima Multitest Swab Specimen Collection Kit.** Immediately place the swab into the transport tube so the black score line is at the top of the tube. Align the score line with the top edge of the tube and carefully break the shaft. The swab will drop to the bottom of the vial. Discard the top portion of the shaft.



Tightly screw the cap onto the tube. When collecting multiple specimens from the same patient, the tube label provides a specimen source field for unique identification for the specimen location.

Hologic provides this collection procedure guide as a general informational tool only; it is not an affirmative instruction or guarantee of performance. It is the sole responsibility of the clinician to read and understand the appropriate package insert and comply with applicable local, state and federal rules and regulations.

hologic.com | diagnostic.solutions@hologic.com | +1.888.484.4747

DS-01552-001 Rev. 003 © 2017 Hologic, Inc. All rights reserved. Hologic, The Science of Sure, Aptima and associated logos are trademarks and/or registered trademarks of Hologic, Inc. and/or its subsidiaries in the United States and/or other countries. This information is intended for medical professionals in the U.S. and other markets and is not intended as a product solicitation or promotion where such activities are prohibited. Because Hologic materials are distributed through websites, eBroadcasts and tradeshows, it is not always possible to control where such materials appear. For specific information on what products are available for sale in a particular country, please contact your Hologic representative or write to diagnostic.solutions@hologic.com.




Aptima® unisex swab kit

Female collection procedure guide


Collection for endocervical swab specimens



Use cleaning swab (white shaft swab with red printing) to remove excess mucus from cervical os and surrounding mucosa. Discard this swab.



Insert collection swab (blue shaft swab with green printing) into endocervical canal. Gently rotate swab clockwise for 10 to 30 seconds to help ensure adequate sampling. Withdraw swab carefully; avoid any contact with vaginal mucosa.



While holding swab in hand, unscrew the tube cap. Do not spill tube contents. **If the tube contents are spilled, discard and replace with a new Aptima unisex swab transport tube.** Carefully break the swab shaft at the score line against the side of the tube. Discard top portion of swab shaft.



Re-cap swab specimen transport tube tightly.

Swab specimen collection guide for:

- *Chlamydia trachomatis* (CT)
- *Neisseria gonorrhoeae* (GC)
- *Trichomonas vaginalis* (TV)

Specimen Transport and Storage

- After collection, transport and store swab in unisex specimen transport tube between 2°C to 30°C until tested.
- Specimens must be assayed with the Aptima assay for CT, GC and /or TV within 60 days of collection.
- If longer storage is needed, freeze between -20°C to -70°C for up to 12 months after collection in the Aptima assay for CT and/or GC. For the Aptima assay for TV, freeze at < -20°C for up to 12 months.




Aptima® unisex swab device

Male collection procedure guide

Collection for male urethral swab specimens

Patient should not have urinated for at least 1 hour prior to specimen collection.



Discard cleaning swab (white shaft with red print on label). The cleaning swab is NOT needed for male specimen collection.

Insert specimen collection swab (blue shaft swab with green printing) 2 cm to 4 cm into urethra. Gently rotate swab clockwise for 2 to 3 seconds in urethra to help ensure adequate sampling. Withdraw swab carefully.



While holding swab in hand, unscrew tube cap. Do not spill tube contents. **If tube contents are spilled, discard and replace with a new Aptima unisex swab transport tube.** Carefully break the swab shaft at the score line against the side of the tube. Discard top portion of swab shaft.



Re-cap swab specimen transport tube tightly.

Swab specimen collection guide for:


- *Chlamydia trachomatis* (CT)
- *Neisseria gonorrhoeae* (GC)

Specimen Transport and Storage

- After collection, transport and store swab in unisex specimen transport tube between 2°C to 30°C until tested.
- Specimens must be assayed with the Aptima assay for CT and/or GC within 60 days of collection.
- If longer storage is needed, freeze between -20°C to -70°C for up to 12 months after collection.

Hologic provides this collection procedure guide as a general informational tool only; it is not an affirmative instruction or guarantee of performance. It is the sole responsibility of the laboratory to read and understand the appropriate package insert and comply with applicable local, state and federal rules and regulations.

hologic.com | info@hologic.com | +1.781.999.7300



Aptima® urine collection kit


Collection procedure guide

Collection for male and female urine specimens

Patient should not have urinated for at least 1 hour prior to specimen collection.



Direct patient to provide **first-catch** urine (approximately 20 to 30 mL of initial urine stream) into urine collection cup free of any preservatives. Collection of larger volumes of urine may result in specimen dilution that may reduce test sensitivity. Female patients should not cleanse labial area prior to providing specimen.



Remove cap from urine specimen transport tube and transfer 2 mL of urine into urine specimen transport tube using the disposable pipette provided. The correct volume of urine has been added when the fluid level is between the black fill lines on urine specimen transport tube label.



Re-cap urine specimen transport tube tightly. This is now known as the “processed urine specimen.”

Urine specimen collection guide for:

- *Chlamydia trachomatis* (CT)
- *Neisseria gonorrhoeae* (GC)
- *Trichomonas vaginalis* (TV) for female only



Specimen transport and storage

- After collection, transport and store processed urine specimens in the Aptima urine specimen transport tube between 2°C to 30°C until tested.
- Processed urine specimens should be assayed with the Aptima assay for CT, GC and/or TV within 30 days of collection.
- If longer storage is needed, freeze between -20°C to -70°C for up to 12 months after collection in the Aptima assay for CT and/or GC. For the Aptima assay for TV, freeze at < -20°C for up to 12 months.
- Urine samples still in primary collection container must be transported to the lab between 2°C to 30°C.
- Transfer urine sample into Aptima urine specimen transport tube within 24 hours of collection.
- Store between 2°C to 30°C and test within 30 days of collection.

Hologic provides this collection procedure guide as a general informational tool only; it is not an affirmative instruction or guarantee of performance. It is the sole responsibility of the laboratory to read and understand the appropriate package insert and comply with applicable local, state and federal rules and regulations.

hologic.com | info@hologic.com | +1.781.999.7300